

# Electrophysiological Indices of Target and Distractor Processing in Visual Search

Clayton Hickey<sup>1,2</sup>, Vincent Di Lollo<sup>2</sup>, and John J. McDonald<sup>2</sup>

## Abstract

■ Attentional selection of a target presented among distractors can be indexed with an event-related potential (ERP) component known as the N2pc. Theoretical interpretation of the N2pc has suggested that it reflects a f

after the onset of a search array and are more pronounced for difficult discrimination tasks than for simple detection tasks and when distractors are near the target rather than far away (Luck, Girelli, et al., 1997).

The appa

button. Four matches were made. In two instances, the initial luminance of the gray patch was  $\sim 2.5 \text{ cd/m}^2$  greater than that of the red, and participants were instructed to decrease the luminance of the gray patch until its brightness matched that of the red patch. In the remaining instances, the initial luminance of the gray patch was  $\sim 2.5 \text{ cd/m}^2$  less than that of the red, and participants were instructed to increase the luminance of the gray patch until its brightness matched that of the red. The numeric values associated with the shade of gray arrived at in each of these matches were mean averaged to define the background luminance used in the experiment.

The experiment itself consisted of 20 blocks of 60 trials, for a total of 1200 trials per participant. All stimuli were presented on a CRT monitor located 60 cm from the observer's eyes. Trials began with the presentation of a fixation point for 1350–1650 msec followed by the presentation of the search array. The array contained two stimuli, a green square that could be rotated 45 to

re-referenced to the algebraic average of the signals recorded at the left and right mastoids. The EEG was amplified with

we

vertical meridian. A clear divergence of the ipsilateral and contralateral waveforms is in evidence in the latency of the P1 (~120–140 msec

Experiment 3 also included displays in which both the target and the distractor were presented on the vertical meridian. The ERP elicited by this nonlateralized display was employed as a baseline to which the lateralized-distractor ERPs could be compared. If the  $P_D$  reflects an ipsilateral negativity, the waveform elicited ipsilateral to an ignored stimulus should be more negative than the waveform elicited by the nonlateralized display. In contrast, if the  $P_D$  reflects a contralateral positivity, the waveform elicited contralateral to the ignored stimulus should be more positive than

## Discussion

Experiment 3 shows that the waveform elicited contralateral to an ignored distractor is, in fact, more positive than that elicited by a nonlateralized display, and thus, demonstrates that the  $P_D$  is a positive ERP component elicited contralateral to the location of an ignored distractor. These results are consistent with the idea that the  $P_D$  reflects a suppressive mechanism that acts on the cortical representation of distractor stimuli.

The experiments reported to this point have been aimed at isolating neural activity tied to distractor processing. What was not addressed was the issue of the corresponding neural activity related to target processing. The electrophysiological activity related to target processing has been the object of earlier investigations that led to the identification of the N2pc (e.g., Eimer, 1996; Luck & Hillyard, 1994a). Those studies employed stimuli displays that were designed to balance sensory energy across the visual hemifields, allowing for lateralized ERP effects to be unambiguously attributed to attention rather than sensory activity. This was done by presenting a salient nontarget item in the hemifield contralateral to the target, such that each visual hemifield contained an equal number of salient and nonsalient visual stimuli. Results from the present experiments, however, suggest a potential drawback to this strategy. Because the N2pc is defined as the difference between signals recorded over the ipsilateral and contralateral h

n

dexRRilite s ref is,

to to

atte

mech

acti



Condition order:  $F(1, 10) = 1.76, p = .214, \text{Condition} \times$   
Condition order:  $F < 1$ ].

#### Electrophysiological Results

Figures 4 and 5 present the ERPs

obtained in the present experiment arose from a display that was not balanced across the visual hemifields. This means that the negativity might reflect a combination of sensory

Figure 6A presents the contralateral-minus-ipsilateral difference waves based on the ERPs presented in Figures 4C and 5C. These difference waves show that the  $N_T$  peaked before the  $P_D$  (250 msec vs. 289 msec, respectively). This latency difference was statistically significant in a RANOVA with a within-participant factor for condition (attend-line vs. attend-square) and a between-participant factor for condition order (attend-line first vs. attend-square first) [Condition:  $F(1, 10) = 5.110$ ,  $p = .045$ ; Condition order:  $F < 1$ ; Condition  $\times$  Condition Order:  $F(1, 10)$ ]

evenly across the visual field would cancel out across the cortical hemispheres.

Hypothetical summation of the  $P_D$  and  $N_T$  components is illustrated in Figure 6. Figure 6A contains two difference waves labeled  $P_D$  and  $N_T$ . The  $P_D$  waveform was calculated by subtracting the contralateral from the ipsilateral waveform in Figure 4C. Similarly, the  $N_T$  waveform was calculated by subtracting the contralateral from the ipsilateral waveform in Figure 5C. The outcome of Experiment 4 suggests that the  $N_T$  represents attentional modulation of neural activity related to target processing—whether excitatory or suppressive. By the same token, the  $P_D$  represents attentional modulation of neural activity related to distractor processing—presumably suppressive. Figure 6B illustrates the abso-

lute algebraic summation of the  $P_D$  and  $N_T$  waveforms.

The waveform in Figure 6B is thus a hypothetical representation of 6009.9626266.054500358.3785207.8364Tm(e)Tj9.96

the

neural activity associated with enhanced target processing, as reported in the animal liter

object features and involved in the processing of target stimuli.

We would like to note that the present results are consistent with what might be expected on the basis of activity at the cellular level. A positive-going effect contralateral to the distractor ( $P_D$ ) and a negative-going effect contralateral to the target ( $N_T$ ) are precisely what would be expected given excitatory postsynaptic potentials (PSPs) in

processes, one tied to the spatial location of distractor stimuli ( $P_D$ ), the other to the spatial location of target stimuli ( $N_T$ ).

## Acknowledgments

We thank Steve Luck and Geoff Woodman for comments on earlier versions of this manuscript. This study was supported in part by grants from the Canadian Institutes of Health Research [C. H.], the Natural Sciences and Engineering Research Council of Canada [V. D. L. and J. J. M.], the Canadian Foundation for Innovation [J. J. M.], and the Canada Research Chairs program [J. J. M.].

Reprint requests should be sent to Clayton Hickey, Department of Cognitive Psychology, Vrije Universiteit Amsterdam, van der Boechorststraat 1, 1081 BT, Amsterdam, The Netherlands, or via e-mail: c.hickey@psy.vu.nl, or to John J. McDonald, Department of Psychology, Simon Fraser University, 8888 University Drive, Burnaby, BC, V5A 1S6, Canada, or via e-mail: jmcd@sfu.ca.

## REFERENCES

- Brown, M. B., & Forsythe, A. B. (1974). Robust tests for equality of variance. *Journal of the American Statistical Association*, *69*, 364–367.
- Cepeda, N. J., Cave, K. R., Bichot, N. P., & Kim, M. S. (1998). Spatial selection via feature-driven inhibition of distractor locations. *Perception & Psychophysics*, *60*, 727–746.
- Chelazzi, L., Miller, E. K., Duncan, J., & Desimone, R. (1993). A neural basis for visual search in inferior temporal cortex. *Nature*, *363*, 345–347.
- Desimone, R., & Duncan, J. (1995). Neural mechanisms of selective visual attention. *Annual Review of Neuroscience*, *18*, 193–222.
- Di Lollo, V., Enns, J. T., & Rensink, R. A. (2000). Competition for consciousness among visual events: The psychophysics of reentrant visual processes. *Journal of Experimental Psychology: General*, *129*, 481–507.
- Eimer, M. (1996). The N2pc component as an indicator of attentional selectivity. *Electroencephalography and Clinical Neurophysiology*, *99*, 225–234.
- Hickey, C.,

Spitzer, H., Desimone, R., & Moran, J. (1988). Increased attention enhances both behavioral and neuronal performance.